



Short communication

Identification of the Changbaishan ‘Millennium’ (B-Tm) eruption deposit in the Lake Suigetsu (SG06) sedimentary archive, Japan: Synchronisation of hemispheric-wide palaeoclimate archives



Danielle McLean^{a,*}, Paul G. Albert^a, Takeshi Nakagawa^{b,c}, Richard A. Staff^a, Takehiko Suzuki^d, Suigetsu 2006 Project Members¹, Victoria C. Smith^a

^a Research Laboratory for Archaeology and the History of Art, University of Oxford, Oxford, OX1 3QY, UK

^b Department of Geography, University of Newcastle, Newcastle Upon Tyne, NE1 7RU, UK

^c Research Centre for Palaeoclimatology, Ritsumeikan University, Kyoto, 603-8577, Japan

^d Department of Geography, Tokyo Metropolitan University, Tokyo, 192-0397, Japan

ARTICLE INFO

Article history:

Received 6 June 2016

Received in revised form

12 August 2016

Accepted 16 August 2016

Available online 25 August 2016

Keywords:

Tephrochronology

Changbaishan ‘Millennium’ eruption

B-Tm tephra

Lake Suigetsu

Bayesian age-modelling

ABSTRACT

The B-Tm tephra, dispersed during the highly explosive Changbaishan ‘Millennium’ eruption (ca. 940–950 CE) and a key marker layer within the Greenland ice cores, has now been identified in the Lake Suigetsu (SG06) sedimentary sequence, central Japan. The major element geochemistry of the volcanic glasses within this tephra layer are compared to a new glass dataset from the distal type-locality (Tomakomai Port, Hokkaido) and other published ‘Millennium’ eruption/B-Tm deposits, to verify this correlation. The discovery of the B-Tm tephra in the Lake Suigetsu record provides, to date, the most southerly identification of this ash and, crucially, the first direct tie-point between this high-resolution, mid-latitude palaeoclimate archive and the Greenland ice cores. These findings present significant encouragement for on-going research into the tephrostratigraphy of East Asia, focusing on the identification of widely-dispersed tephra layers which can facilitate the synchronisation of disparate palaeoclimate archives and thus enable the assessment of spatio-temporal variations in past climatic change.

© 2016 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Our understanding of past climate systems and environmental change is formed through the comparison of numerous multiproxy palaeoenvironmental archives. Volcanic ash (tephra) horizons have proved extremely valuable for such studies as they allow disparate sedimentary records to be precisely correlated and dated (e.g. Lane et al., 2014; Albert et al., 2015; Lowe et al., 2015; Abbott et al., 2016). Such isochronous tie-points are essential to assessing the spatio-temporal variability of major climatic events and to assess abrupt and short-lived fluctuations, controlled by driving systems such as the East Asian monsoon (EAM). The EAM is of major socioeconomic importance for the ~30% of the global population that reside in East Asia, yet its driving complexities are poorly understood and climatically modelled, partly due to limited

long-term observations (Liu and Ding, 1998; Cook et al., 2010; Ha et al., 2012). In addition, high-resolution palaeoenvironmental archives, such as the sediments of Lake Suigetsu in central Japan (Nakagawa et al., 2003, 2005), indicate that the magnitude and timing of major palaeoclimatic events do not necessarily occur synchronously with those reported in the polar and high-latitude regions. The identification of more widely-dispersed tephra layers, particularly connecting to key climate stratotypes including the Greenland ice cores, are fundamental to assessing the interplay between these regions and the forcing mechanisms of palaeomonsoon variability.

Here we outline the recent discovery of the Baegdusan-Tomakomai (B-Tm) tephra layer within the sediments of Lake Suigetsu, and thus the first direct tephra tie-point between this high-resolution, mid-latitude lacustrine palaeoclimate archive and the Greenland ice cores. This hemispheric-wide tephrostratigraphic marker is precisely dated using the Greenland (NS1-2011) chronology (Sigl et al., 2015) and therefore contributes a robust constraint to the Late Holocene sedimentation at Lake Suigetsu.

* Corresponding author.

E-mail address: danielle.mclean@jesus.ox.ac.uk (D. McLean).

¹ www.suigetsu.org.

2. Background

2.1. The B-Tm tephra

The B-Tm tephra was dispersed during the highly explosive 'Millennium' eruption of Changbaishan (also referred to as Tianchi or Baitoushan) volcano, situated on the border between China and North Korea (128°03'E, 41°00'N) (Machida et al., 1990) (Fig. 1). This was one of the most violent eruptions of the past 2000 years, with a volcanic explosivity index (VEI) (Newhall and Self, 1982) of 7 (i.e. comparable to the 1815 CE eruption of Tambora, Indonesia).

The eruption had two explosive phases, with the initial main phase (ca. 95% by volume) associated with a ca. 25 km-high Plinian column, producing a widespread pumice fall unit (Machida et al., 1990; Horn and Schmincke, 2000). This fall unit is overlain by partially-welded pyroclastic density current (PDC) deposits attributed to the partial collapse of the Plinian column and are comenditic in composition. Trachytic magma was erupted in a late phase of the eruption, forming moderately welded PDC units, that post-date the comenditic fall and PDC deposits (Horn and Schmincke, 2000).

The B-Tm tephra was named and characterised (using glass

refractive indices and major element compositions) at a distal type-locality in Tomakomai Port, Hokkaido (42°39'18"N, 141°41'53"E; 53 km SSE of Sapporo City), where it was identified above the Tarumai-c (ca. 50 BCE) and below the Tarumai-b (1667 CE) tephra layers (from the nearby Tarumae volcano) (Machida and Arai, 1983). The B-Tm tephra has since been identified in numerous marine, lacustrine and archaeological sequences across northern Japan, northeast China and coastal regions of Russia (Fig. 1). Moreover, a recent study found B-Tm glass shards in the Greenland ice cores, which provided a precise ice core-derived eruption age of 941 CE \pm 1 (Sun et al., 2014b). ^{10}Be and ^{14}C tree-ring markers have since shown that there is a seven-year offset in the Greenland ice core (GICC05) timescale, so this age for the B-Tm eruption has been corrected to 946–947 CE (NS1-2011; Sigl et al., 2015).

2.2. Lake Suigetsu (SG06) archive

The Lake Suigetsu (Fig. 1; 35°35'0"N; 135°53'0"E) SG06 sediment core provides a high-resolution palaeoenvironmental record from Honshu Island, central Japan (e.g. Tyler et al., 2010; Kossler et al., 2011; Saito-Kato et al., 2013; Schlolaut et al., 2014). The




Fig. 1. Location of Changbaishan volcano (China/North Korea border), Lake Suigetsu (central Japan) and other sites that preserve the B-Tm tephra. Pie charts indicate the proportion of trachytic (black) and comenditic (white) glasses in sites where there is sufficient major element glass data published to form a meaningful assessment of correlation ($n > 10$; shown in the middle of each chart; n = number of shards). Squares represent sites that report the B-Tm tephra, based on physical characteristics of the unit, but with limited or no glass geochemical data. Dispersal boundaries (dashed line) for the B-Tm as previously mapped by Machida and Arai (1983). References: (a) this study; (b) Sun et al. (2015); (c) Furuta et al. (1986); Machida and Arai (1983); (d) Chen et al. (2016); (e) Wada et al. (2001); (f) Nakamura (2016); (g) this study; (h) Hughes et al. (2013); (i) Nakagawa et al. (2002); (j) Nanayama et al. (2003); (k) Fukusawa et al. (1998); (l) Okuno et al. (2011); (m) Sun et al. (2014b); (n) Coulter et al. (2012).

73.19 m-long composite core was extracted as part of the ‘Suigetsu Varves 2006’ project, and provides a continuous record of sedimentation spanning the last ca. 150 ka (Nakagawa et al., 2012). The sequence is annually laminated (varved) between ca.10 and 70 ka and has been extensively radiocarbon (^{14}C) dated and varve counted to generate a high-resolution chronology (Staff et al., 2011; Bronk Ramsey et al., 2012; Marshall et al., 2012; Schlolaut et al., 2012) for this palaeoenvironmental record.

Hitherto, 31 visible tephra layers from SG06 have been identified and characterised (using major element geochemical analysis),

and provide a detailed record of Japanese and Korean eruption events (Smith et al., 2013). Due to the prevailing westerlies, tephra is largely dispersed northeast across this region. As Changbaishan is located northwest of Lake Suigetsu, coupled with the strong easterly dispersal of the eruptive plume implied from ash thicknesses (Fig. 1), the presence of a B-Tm ash layer in the sediments was previously considered unlikely. However, given its widespread distribution and recent ultra-distal identification in the Greenland ice cores (Fig. 1), the Lake Suigetsu sediments were carefully re-examined in order to attempt to identify this key tephra marker.

Table 1

Results of the WDS-EPMA normalised glass compositions of the SG06-0226 tephra layer and of the B-Tm distal type-site deposits at Tomakomai Port.

Sample	SiO ₂	TiO ₂	Al ₂ O ₃	FeO	CaO	Na ₂ O	K ₂ O	Cl	Analytical total
SG06-0226	74.96	0.18	10.23	3.99	0.22	5.34	4.46	0.51	97.99
	74.95	0.26	10.14	3.97	0.24	5.43	4.40	0.52	97.90
	74.83	0.26	10.30	4.20	0.23	5.29	4.41	0.48	98.03
	75.14	0.19	10.17	4.02	0.23	5.34	4.35	0.46	97.31
	75.06	0.18	10.18	4.02	0.22	5.45	4.33	0.50	97.38
	74.89	0.23	10.32	3.97	0.23	5.51	4.21	0.49	97.36
	75.17	0.24	10.18	3.95	0.23	5.19	4.43	0.48	96.97
	74.84	0.21	10.22	3.92	0.18	5.71	4.40	0.46	97.35
	74.57	0.19	10.28	4.27	0.18	5.51	4.44	0.50	97.59
	74.85	0.19	10.53	3.90	0.21	5.33	4.38	0.47	97.15
	74.84	0.19	10.22	4.05	0.21	5.53	4.31	0.50	96.91
	74.65	0.28	10.28	4.19	0.20	5.42	4.27	0.52	96.88
	75.20	0.23	10.19	4.19	0.24	5.12	4.22	0.49	96.17
	74.87	0.29	10.39	3.98	0.29	5.19	4.35	0.50	96.56
	74.67	0.23	10.31	4.10	0.24	5.49	4.43	0.48	96.72
	74.96	0.19	10.22	3.99	0.19	5.48	4.25	0.50	96.33
	74.89	0.23	10.28	4.06	0.22	5.42	4.30	0.48	96.39
	74.80	0.24	10.30	4.27	0.23	5.33	4.27	0.50	96.42
	75.04	0.17	10.19	4.20	0.21	5.20	4.41	0.46	95.89
	75.49	0.20	10.20	3.72	0.22	5.09	4.44	0.48	95.12
	74.80	0.23	10.39	3.91	0.24	5.46	4.27	0.52	95.83
	74.94	0.25	10.25	4.28	0.21	5.26	4.21	0.52	95.38
	75.08	0.20	10.24	4.05	0.19	5.27	4.45	0.44	94.53
	75.06	0.20	10.25	4.20	0.21	5.17	4.42	0.47	94.55
	74.61	0.27	10.43	3.73	0.23	5.62	4.44	0.56	94.93
	74.55	0.26	10.30	4.13	0.22	5.36	4.50	0.51	94.98
	74.61	0.23	10.49	4.06	0.23	5.16	4.55	0.54	94.05
	74.88	0.18	10.23	4.02	0.23	5.45	4.48	0.51	93.48
	74.73	0.20	10.16	4.17	0.23	5.34	4.48	0.52	93.22
	65.14	0.57	16.10	4.55	1.48	5.41	6.15	0.10	96.87
Average (n = 30)	74.57	0.23	10.47	4.07	0.26	5.36	4.43	0.48	
1σ	1.79	0.07	1.07	0.17	0.23	0.15	0.34	0.08	
SG06-0226 outliers	77.50	0.18	12.61	1.22	1.11	3.75	3.27	0.12	94.00
	73.86	0.48	12.97	2.45	1.95	3.08	4.35	0.14	98.28
Tomakomai Port	74.80	0.18	10.46	3.98	0.25	5.36	4.31	0.59	93.49
	75.27	0.22	10.65	3.92	0.25	4.80	4.31	0.52	96.91
	74.51	0.23	10.65	4.08	0.31	5.29	4.36	0.47	97.88
	75.13	0.18	10.51	4.08	0.27	4.98	4.26	0.47	99.41
	67.33	0.44	15.00	4.65	1.20	5.90	4.98	0.19	98.78
	74.88	0.21	10.51	3.87	0.25	5.28	4.37	0.50	97.90
	74.97	0.23	10.58	3.98	0.23	5.25	4.19	0.48	97.73
	74.41	0.25	10.86	3.85	0.30	5.07	4.62	0.52	97.82
	74.47	0.23	11.08	3.90	0.33	5.09	4.31	0.50	98.54
	75.08	0.20	10.59	3.97	0.23	5.10	4.21	0.55	96.35
	74.67	0.17	10.35	4.26	0.32	5.29	4.29	0.59	95.24
	74.10	0.28	10.90	4.14	0.28	5.34	4.44	0.47	98.96
	74.53	0.26	10.72	3.99	0.22	5.20	4.49	0.49	97.22
	74.46	0.25	10.54	4.11	0.30	5.31	4.41	0.56	93.69
	67.05	0.36	15.02	4.62	1.05	6.04	5.43	0.17	98.60
	69.91	0.30	13.66	4.26	0.59	5.81	5.12	0.31	94.54
	73.95	0.19	10.68	4.09	0.29	5.82	4.37	0.51	97.36
	75.01	0.18	10.40	4.05	0.26	5.08	4.41	0.48	96.71
	74.68	0.20	10.42	4.11	0.24	5.27	4.51	0.46	97.60
	74.62	0.20	10.35	3.96	0.24	5.51	4.50	0.51	96.90
	74.20	0.25	10.71	4.12	0.29	5.44	4.49	0.47	97.79
Average (n = 21)	73.72	0.24	11.17	4.09	0.37	5.34	4.49	0.47	
1σ	2.43	0.07	1.45	0.21	0.26	0.32	0.31	0.11	

The data have been filtered to remove analyses of a <93% analytical total; MgO, MnO and P₂O₃ values are below the statistical detection threshold and have been removed (see supplementary material).

3. Materials and methods

3.1. Sampling and laboratory procedures

A new, thin ash layer (<1 mm) was observed within overlapping core sections of the SG06 sequence at a composite depth (CD) of 225.5 cm (the CD was determined using the 20 April 2015 version of the correlation model; see Nakagawa et al., 2012 for more details). Herein, this tephra is labelled as SG06-0226. A subsample was extracted from core section D-01 at 104.7cm depth and wet sieved using distilled water through a 25 μm mesh. Microscopic examination identified an abundant concentration of colourless glass shards with a distinct, highly vesicular morphology. These were handpicked using a micromanipulator (Lane et al., 2014) and mounted in epoxy resin for geochemical analysis. To confirm the stratigraphic positioning of the peak, cryptotephra analysis was performed in sediments above and below the layer. A sample of the B-Tm tephra from its type-locality at Tomakomai Port was also prepared for geochemical analysis to verify the correlation.

3.2. Compositional analysis

Major and minor element glass compositions were measured using a JEOL-8600 wavelength-dispersive electron microprobe (WDS-EPMA) at the Research Laboratory for Archaeology and History of Art, University of Oxford. All analyses were performed

using an accelerating voltage of 15 kV, beam current of 6 nA and 10 μm -diameter beam. The electron microprobe was calibrated using a suite of mineral standards, and the PAP absorption correction method was used for quantification. The precision and accuracy of these analyses were assessed using the MPI-DING reference glasses (Jochum et al., 2006), which were run with our samples. All data comparisons are based on normalised glass compositions.

4. Results and discussion

4.1. Geochemical results

The geochemical analyses of glass shards from the SG06-0226 layer and from the B-Tm type-locality (Tomakomai Port) are shown in Table 1 and selected bi-plots in Fig. 2. Volcanic glasses erupted from the Changbaishan volcano can be easily discriminated from those originating from volcanic centres in Russia (Kamchatka), South Korea and along the Japanese arc (Fig. 2a). The eruption that produced the B-Tm tephra tapped both comenditic and trachytic magma batches, with particularly alkali-enriched compositions. The SG06-0226 dataset includes two outliers, which have the same composition as other shards present in low concentrations throughout some other sections of the core and are considered to be background material. These outlier glass compositions are chemically distinct from the B-Tm tephra and clearly erupted from a Japanese volcano (Fig. 2a), so are therefore not discussed further.




Fig. 2. Major element glass compositions of the SG06-0226 tephra (red circles) and the B-Tm type-locality from Tomakomai Port (green squares) (a) Total alkalis versus silica plot (TAS; Le Bas et al., 1986) showing the glass composition of Changbaishan tephra relative to those from Ulleung Island (Smith et al., 2013), volcanoes in the Japanese arc (Smith et al., 2013), and Kamchatka (Kyle et al., 2011); (b) peralkaline classification diagram (McDonald, 1974); (c) the composition of glass shards from proximal Changbaishan 'Millennium' eruption deposits (Guo et al., 2002; Zou et al., 2010; Sun et al., 2014b, 2015; Chen et al., 2016) and composite published distal B-Tm data (Furuta et al., 1986; Okuno et al., 2011; Coulter et al., 2012; Hughes et al., 2013; Sun et al., 2014b, 2015; Chen et al., 2016); (d) the glass composition of distal B-Tm units compared to that of the SG06-0226 glasses. Error bars represent 2 x standard deviation of repeat analyses of the Sth6/80-G MPI-DING standard glass. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

The SG06-0226 glasses are compositionally heterogeneous, with SiO₂ ranging from 65.14 to 75.49 wt%, CaO from 0.18 to 1.48 wt%, FeOt (all Fe as FeO) from 3.72 to 4.55 wt% and K₂O from 4.21 to 6.15 wt% (n = 30). The deposit is dominated by a comenditic glass component (Fig. 2), with a single glass shard showing a trachytic composition. The comenditic glasses contain a lower Al₂O₃ (ca. 10 wt%) and FeOt (ca. 4.0 wt%) content compared to the single trachytic glass shard (Fig. 2b). The glass compositions of SG06-0226 are consistent with the new glass data from the B-Tm type-locality at Tomakomai Port, confirming they are the same tephra. The new Tomakomai Port dataset also contains compositions that lie on a mixing line between the comenditic and trachytic end members of the SG06-0226 tephra, which suggests that the two distinct magma batches were in contact before the eruption.

The geochemistry of the SG06-0226 and Tomakomai Port glasses are also consistent with published data for proximal Changbaishan 'Millennium' eruption deposits (Fig. 2c) and other distal/ultra-distal B-Tm glass compositions, including those preserved in the Greenland ice cores (Fig. 2d). The glass compositions of the tephra at these distal sites provide information on the spatial distribution of the different phases of the Changbaishan 'Millennium' eruption (Fig. 1). Given that the trachytic compositions were erupted in the late PDC phase of the eruption, the dispersal of glass with such compositions to the distal sites must have been via the plume that lofted off the top of the PDCs, i.e., a co-PDC plume. The compositions from the sites directly to the east of Changbaishan are predominately comenditic (ca. 50–80%), and therefore mostly associated with the Plinian column or the co-PDC plume associated with the PDC generated during the collapse of the Plinian column, or both. These sites also have a trachytic component (ca. 15–50%) implying that the dispersal axis for the late co-PDC plume was also to the east. The compositions of glass shards from sites northwest of the vent are mostly trachytic, implying that these glasses were deposited from the co-PDC plume. Glass shards in SG06-0226 are almost entirely comenditic in composition and associated with the Plinian column and the co-PDC associated with its collapse, and indicate that the late phase trachytic co-PDC plume did not disperse tephra towards the south. The trachytic compositions are more dominant in the Greenland ice cores implying that the tephra was mostly transported ultra-distally (~7000 km from vent) via the co-PDC plume. This inference is consistent with some studies of other large explosive eruptions that show that the ultra-distal transport of tephra is largely associated with co-PDC plumes (e.g., Campanian Ignimbrite eruption, Smith et al., 2016; Younger Toba Tuff, Costa et al., 2014).

4.2. Age-depth modelling

The age model for the SG06 sedimentary sequence (Staff et al., 2011) was constructed using a Bayesian Poisson-process 'P_Sequence' depositional model in OxCal (ver. 4.2; Bronk Ramsey, 2008, 2016), and the IntCal13 calibration curve (Reimer et al., 2013) (Fig. 3). This model includes 124 AMS ¹⁴C dates obtained from terrestrial plant macrofossils from the non-varved Holocene part of the core (Staff et al., 2011). An age of 880–956 cal. CE (95.4% confidence) was obtained for the SG06-0226 tephra, consistent with other published dates for the 'Millennium' eruption (see Sun et al., 2014a). The lack of varved sedimentation throughout the late Holocene means that this SG06-derived age is not as precise as those reported elsewhere. We are, however, able to import the NS1–2011 age of 947 CE ± 1 (Sigl et al., 2015) into the Lake Suigetsu age model, thus further constraining the chronology of the upper part of SG06. This imported ice-core-based age will significantly improve the chronological analysis of proxy palaeoenvironmental data and any subsequent cryptotephra layers identified in the Suigetsu sediments within this timeframe.




Fig. 3. The age-depth model for the upper 5 m of the SG06 core (after Staff et al., 2011). The blue envelope represents the 95.4% probability range. Inset: posterior probability density function (PDF) generated for the deposition of the SG06-0226 tephra, located at 219.1 cm event-free CD (determined using 29 Jan. 2011 version of the event-free depth model). The bars underlying the PDF represent the 68.2%, 95.4% and 99.7% probability ranges, respectively. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

5. Implications

5.1. Palaeoenvironmental potential for the B-Tm tephra

The stratigraphic positioning of the B-Tm tephra holds significant potential to help constrain late Holocene climatic variations, such as the onset of the Medieval Warm Period (ca. 900–1300 CE). Although this warming episode has been identified in numerous global paleoclimate proxy records, its spatio-temporal variability and regional magnitude is intensely debated (Broecker, 2001; Hunt, 2006). Once high-resolution palaeoenvironmental proxy data from Lake Suigetsu are obtained from this interval, the dynamics of such rapid and short-lived climatic episodes can be evaluated. Our discovery may also encourage others to look for the B-Tm tephra in other key palaeoenvironmental records, including sites further south.

5.2. Future potential

This study provides the first direct tephra tie-point between a high-resolution, mid-latitude palaeoclimate archive (Lake Suigetsu sediments) and the Greenland ice cores, a major goal of research communities examining palaeoenvironmental change (e.g.

INTIMATE; Lowe et al., 2015; Blockley et al., 2014). This discovery therefore provides significant encouragement for the on-going research into refining the tephrostratigraphy of East Asia, research which aims to establish more hemispheric-wide tephra tie-points suitable for the synchronisation of key palaeoclimate archives.

Acknowledgements

This work was supported by the UK Natural Environmental Research Council (NERC; grants: NE/D00289/1 and NE/F003048/1), the Deutsche Forschungsgemeinschaft (DFG; grant: TA-540/3-1) and MEXT-Japan KAKENHI (grant: 21101002 and 15H02143), which funded the Suigetsu Varves 2006 Project. DM is funded by NERC (grant: NE/L002612/1) and part of the Environmental Research Doctoral Training Program at the University of Oxford. PGA and RAS are supported by Early Career Fellowships from the Leverhulme Trust (grant: ECF-2014-438 and ECF-2015-396). The authors would like to thank: Prof. Hiroshi Machida for providing the B-Tm type-locality sample from Tomakomai Port; and David Lowe and an anonymous reviewer for their comments.

Appendix A. Supplementary data

Supplementary data related to this article can be found at <http://dx.doi.org/10.1016/j.quascirev.2016.08.022>.

References

- Abbott, P.M., Bourne, A.J., Purcell, C.S., Davies, S.M., Scourse, J.D., Pearce, N.J.G., 2016. Last glacial period cryptotephra deposits in an eastern North Atlantic marine sequence: exploring linkages to the Greenland ice-cores. *Quat. Geochronol.* 31, 62–76.
- Albert, P.G., Hardiman, M., Keller, J., Tomlinson, E.L., Smith, V.C., Bourne, A.J., Wulf, S., Zanchetta, G., Sulpizio, R., Müller, U.C., Pross, J., Ottoloni, L., Matthews, I.P., Blockley, S.P.E., Menzies, M.A., 2015. Revisiting the Y-3 tephrostratigraphic marker: a new diagnostic glass geochemistry, age estimate, and details on its climatostratigraphical context. *Quat. Sci. Rev.* 118, 105–121.
- Blockley, S.P.E., Bourne, A.J., Brauer, A., Davies, S.M., Hardiman, M., Harding, P.R., Lane, C.S., MacLeod, A., Matthews, I.P., Pyne-O'Donnell, S.D.F., Rasmussen, S.O., Wulf, S., Zanchetta, G., 2014. Tephrochronology and the extended INTIMATE (integration of ice-Core, marine and terrestrial records) event stratigraphy 8–128 ka b2k. *Quat. Sci. Rev.* 106, 88–100.
- Broecker, W.S., 2001. Was the medieval Warm period global? *Science* 291, 1497–1499.
- Bronk Ramsey, C., 2008. Deposition models for chronological records. *Quat. Sci. Rev.* 27, 42–60.
- Bronk Ramsey, C., 2016. OxCal Project, Version 4.2. Retrieved April 2016. <https://c14.arch.ox.ac.uk/oxcal/OxCal.html>.
- Bronk Ramsey, C., Staff, R.A., Bryant, C.L., Brock, F., Kitagawa, H., Van Der Plicht, J., Schlögl, G., Marshall, M.H., Brauer, A., Lamb, H.F., Payne, R.L., Tarasov, P.E., Haraguchi, T., Gotanda, K., Yonenobu, H., Yokoyama, Y., Tada, R., Nakagawa, T., 2012. A complete terrestrial radiocarbon record for 11.2 to 52.8 kyr BP. *Science* 338, 370–374.
- Chen, X.-Y., Blockley, S.P.E., Tarasov, P.E., Xu, Y.-G., McLean, D., Tomlinson, E., Albert, P.G., Liu, J., Muller, S., Wagner, M., Menzies, M.A., 2016. Clarifying the distal to proximal tephrochronology of the Millennium Eruption, Changbaishan, northeast China. *Quat. Geochronol.* 33, 61–75.
- Cook, E.R., Anchukaitis, K.J., Buckley, B.M., D'Arrigo, R.D., Jacoby, G.C., Wright, W.E., 2010. Asian monsoon failure and megadrought during the last Millennium. *Science* 328, 486–489.
- Costa, A., Smith, V.C., Macedonio, G., Matthews, N.E., 2014. The magnitude and impact of the Youngest Toba Tuff super-eruption. *Front. Earth Sci.* 2, 16.
- Coulter, S.E., Pilcher, J.R., Plunkett, G., Baillie, M., Hall, V.A., Steffensen, J.P., Vinther, M.B., Clausen, H.B., Johnsen, S.J., 2012. Holocene tephtras highlight complexity of volcanic signals in Greenland ice cores. *J. Geophys. Res.* 117.
- Fukushima, H., Tsukamoto, S., Tsukamoto, H., Ikeda, M., Okamura, M., Matsuoka, M., 1998. Falling age of Baegdusan-Tomakomai tephra (B-Tm) estimated by using non-glacial varves. *Laguna* 5, 55–62.
- Furuta, T., Fujioka, K., Arai, F., 1986. Widespread submarine tephtras around Japan: Petrographic and chemical properties. *Mar. Geol.* 72, 125–142.
- Guo, Z., Liu, J., Sui, S., Liu, Q., He, H., Ni, Y., 2002. The mass estimation of volatile emission during 1199/1200 AD eruption of Baitoushan volcano and its significance. *Sci. Chin. Earth Sci.* 46, 530–539.
- Ha, K.J., Heo, K.Y., Lee, S.S., Yun, K.S., Jhun, J.G., 2012. Variability in the east asian monsoon: a review. *Meteorol. Appl.* 19, 200–215.
- Horn, S., Schmincke, H.U., 2000. Volatile emission during the eruption of Baitoushan Volcano (China/North Korea) ca. 969 AD. *Bull. Volcanol.* 61, 537–555.
- Hughes, P.D.M., Mallon, G., Brown, A., Essex, H.J., Stanford, J.D., Hotes, S., 2013. The impact of high tephra loading on late-Holocene carbon accumulation and vegetation succession in peatland communities. *Quat. Sci. Rev.* 67, 160–175.
- Hunt, B.G., 2006. The medieval Warm period, the little ice age and simulated climatic variability. *Clim. Dyn.* 27, 677–694.
- Jochum, K.P., Stoll, B., Herwig, K., Willbold, M., Hofmann, A.W., Amini, M., et al., 2006. MPI-DING reference glasses for in situ microanalysis: new reference values for element concentrations and isotope ratios. *Geochemistry. Geophys. Geosystems* 7, 2.
- Kossler, A., Tarasov, P., Schlögl, G., Nakagawa, T., Marshall, M., Brauer, A., Staff, R., Ramsey, C.B., Bryant, C., Lamb, H., Damske, D., Gotanda, K., Haraguchi, T., Yokoyama, Y., Tada, R., Suigetsu, 2006 project members, 2011. Onset and termination of the late-glacial climate reversal in the high-resolution diatom and sedimentary records from the annually laminated SG06 core from Lake Suigetsu, Japan. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 306, 103–115.
- Kyle, P.R., Ponomareva, V.V., Rourke Schlup, R., 2011. Geochemical characterization of marker tephra layers from major Holocene eruptions, Kamchatka Peninsula, Russia. *Int. Geol. Rev.* 53, 1059–1097.
- Lane, C.S., Cullen, V.L., White, D., Bramham-Law, C.W.F., Smith, V.C., 2014. Cryptotephra as a dating and correlation tool in archaeology. *J. Archaeol. Sci.* 42, 42–50.
- Le Bas, M.J., Le Maitre, R.W., Streckeisen, A., Zanettin, B., 1986. A chemical classification of volcanic rocks based on the total alkali-silica diagram. *J. Petrol.* 27, 745–750.
- Liu, T., Ding, Z., 1998. Chinese loess and the palaeomonsoon. *Annu. Rev. Earth Planet. Sci.* 26, 111–145.
- Lowe, J.J., Ramsey, C.B., Housley, R., Lane, C., Tomlinson, E.L., RESET Team and RESET Associates, 2015. The RESET project: Constructing a European tephra lattice for refine synchronisation of environmental and archaeological events during the last c. 100 ka. *Quat. Sci. Rev.* 27, 1–17.
- Machida, H., Arai, F., 1983. Extensive ash falls in and around the Sea of Japan from large late Quaternary eruptions. *J. Volcanol. Geotherm. Res.* 18, 151–164.
- Machida, H., Moriwaki, H., Zhao, D., 1990. The recent major eruption of Changbai volcano and its environmental effects. *Geogr. Rep. Tokyo Metropol. Univ.* 25, 1–20.
- Marshall, M., Schlögl, G., Nakagawa, T., Lamb, H., Brauer, A., Staff, R., Ramsey, C.B., Tarasov, P., Gotanda, K., Haraguchi, T., Yokoyama, Y., Yonenobu, H., Tada, R., Suigetsu 2006 Project members, 2012. A novel approach to varve counting using μ XRF and X-radiography in combination with thin-section microscopy, applied to the Late Glacial chronology from Lake Suigetsu, Japan. *Quat. Geochronol.* 13, 70–80.
- McDonald, R., 1974. Nomenclature and petrochemistry of the peralkaline oversaturated extrusive rocks. *Bull. Volcanol.* 38, 498–516.
- Nakagawa, M., Ishizuka, Y., Kudo, T., Yoshimoto, M., Hirose, W., Ishizaki, Y., Gouchi, N., Katsui, Y., Solovyov, A.W., Steinberg, G.S., Abdurakhmanov, A.I., 2002. Tyatya volcano, southwestern Kuril arc: recent eruptive activity inferred from widespread tephra. *Isl. Arc* 11, 236–254.
- Nakagawa, T., Kitagawa, H., Yasuda, Y., Tarasov, P.E., Nishida, K., Gotanda, K., Sawai, Y., Yangtze River and Civilization Program members, 2003. Asynchronous Climate Changes in the North Atlantic and Japan During the Last Termination. *Science* 299, 688–691.
- Nakagawa, T., Kitagawa, H., Yasuda, Y., Tarasov, P.E., Gotanda, K., Sawai, Y., 2005. Pollen/event stratigraphy of the varved sediment of Lake Suigetsu, central Japan from 15,701 to 10,217 SG v. BP (Suigetsu varve years before present): description, interpretation, and correlation with other regions. *Quat. Sci. Rev.* 24, 1691–1701.
- Nakagawa, T., Gotanda, K., Haraguchi, T., Danhara, T., Yonenobu, H., Brauer, A., Yokoyama, Y., Tada, R., Takemura, K., Staff, R.A., Payne, R., Bronk Ramsey, C., Bryant, C., Brock, F., Schlögl, G., Marshall, M., Tarasov, P., Lamb, H., Suigetsu 2006 Project Members, 2012. SG06, a fully continuous and varved sediment core from Lake Suigetsu, Japan: stratigraphy and potential for improving the radiocarbon calibration model and understanding of late Quaternary climate changes. *Quat. Sci. Rev.* 36, 164–176.
- Nakamura, Y., 2016. Stratigraphy, distribution, and petrographic properties of Holocene tephtras in Hokkaido, northern Japan. *Quat. Int.* 397, 52–62.
- Nanayama, F., Satake, K., Furukawa, R., Shimokawa, K., Atwater, B.F., Shigeno, K., Yamaki, S., 2003. Unusually large earthquakes inferred from tsunami deposits along the Kuril trench. *Nature* 424, 660–663.
- Newhall, C.G., Self, S., 1982. The volcanic explosivity index (VEI) an estimate of explosive magnitude for historical volcanism. *J. Geophys. Res.* 87, 1231–1238.
- Okuno, M., Torii, M., Yamada, K., Shinozuka, Y., Danhara, T., Gotanda, K., Yonenobu, H., Yasuda, Y., 2011. Widespread tephtras in sediments from Lake Ichi-no-Megata in northern Japan: their description, correlation and significance. *Quat. Int.* 246, 270–277.
- Reimer, P.J., Bard, E., Bayliss, A., Beck, J.W., Blackwell, P.G., Bronk Ramsey, C., Buck, C.E., Cheng, H., Edwards, L.R., Friedrich, M., Grootes, P.M., Guilderson, T.P., Hafflidason, H., Hajdas, I., Hatté, C., Heaton, T.J., Hoffmann, D.L., Hogg, A.G., Hughes, K.A., Kaiser, K.F., Kromer, B., Manning, S.W., Niu, M., Reimer, R.W., Richards, D.A., Scott, E.M., Southon, J.R., Staff, R.A., Turney, C.S.M., van der Plicht, J., 2013. IntCal13 and Marine13 radiocarbon age calibration curves 0–50,000 years cal BP. *Radiocarbon* 55, 1869–1887.
- Saito-Kato, M., Yamada, K., Staff, R.A., Nakagawa, T., Yonenobu, H., Haraguchi, T., Takemura, K., Bronk Ramsey, C., 2013. An assessment of the magnitude of the

- AD 1586 Tensho tsunami inferred from Lake Suigetsu sediment cores. *J. Geogr. (Chigaku Zasshi)* 122, 493–501 (in Japanese, with English abstract).
- Schlolaut, G., Marshall, M.H., Brauer, A., Nakagawa, T., Lamb, H.F., Staff, R.A., Bronk Ramsey, C., Bryant, C.L., Brock, F., Kossler, A., Tarasov, P.E., Yokoyama, Y., Tada, R., Haraguchi, T., Suigetsu 2006 project members, 2012. An automated method for varve interpolation and its application to the Late Glacial chronology from Lake Suigetsu, Japan. *Quat. Geochronol.* 13, 52–69.
- Schlolaut, G., Brauer, A., Marshall, M.H., Nakagawa, T., Staff, R.A., Ramsey, C.B., Lamb, H.F., Bryant, C.L., Naumann, R., Dulski, P., Brock, F., Yokoyama, Y., Tada, R., Haraguchi, T., Suigetsu 2006 project members, 2014. Event layers in the Japanese Lake Suigetsu 'SG06' sediment core: description, interpretation and climatic implications. *Quat. Sci. Rev.* 83, 157–170.
- Sigl, M., Winstrup, M., McConnell, J.R., Welten, K.C., Plunkett, G., Ludlow, F., Büntgen, U., Caffee, M., Chellman, N., Dahl-Jensen, D., Fischer, H., Kipfstuhl, S., Kostick, C., Maselli, O.J., Mekhaldi, F., Mulvaney, R., Muscheler, R., Pasteris, D.R., Pilcher, J.R., Salzer, M., Schüpbach, S., Steffensen, J.P., Vinther, B.M., Woodruff, T.E., 2015. Timing and climate forcing of volcanic eruptions for the past 2,500 years. *Nature* 523, 543–549.
- Smith, V.C., Staff, R.A., Blockley, S.P.E., Bronk Ramsey, C., Nakagawa, T., Mark, D.F., Takemura, K., Danhara, T., Suigetsu 2006 Project Members, 2013. Identification and correlation of visible tephras in the Lake Suigetsu SG06 sedimentary archive, Japan: chronostratigraphic markers for synchronising of east Asian/west Pacific palaeoclimatic records across the last 150 ka. *Quat. Sci. Rev.* 61, 121–137.
- Smith, V.C., Isaia, R., Engwell, S.L., Albert, P.G., 2016. Tephra dispersal during the Campanian Ignimbrite (Italy) eruption: implications for ultra-distal ash transport during the large caldera-forming eruption. *Bull. Volcanol.* 78, 45–60.
- Staff, R.A., Bronk Ramsey, C., Bryant, C.L., Brock, F., Payne, R.L., Schlolaut, G., Marshall, M.H., Brauer, A., Lamb, H.F., Tarasov, P., Yokoyama, Y., Haraguchi, T., Gotanda, K., Yonenobu, H., Nakagawa, T., Suigetsu 2006 project members, 2011. New ¹⁴C determinations from Lake Suigetsu, Japan: 12,000 to 0 cal bp. *Radiocarbon* 53, 511–528.
- Sun, C., You, H., Liu, J., Li, X., Gao, J., Chen, S., 2014a. Distribution, geochemistry and age of the Millennium eruptives of Changbaishan volcano, Northeast China—A review. *Front. Earth Sci.* 8, 216–230.
- Sun, C., Plunkett, G., Liu, J., Zhao, H., Sigl, M., McConnell, J.R., Pilcher, J.R., Vinther, B., Steffensen, J.P., Hall, V., 2014b. Ash from Changbaishan Millennium eruption recorded in Greenland ice: implications for determining the eruptions timing and impact. *Geophys. Res. Lett.* 41, 694–701.
- Sun, C., You, H., He, H., Zhang, L., Gao, J., Guo, W., Chen, S., Mao, Q., Liu, Q., Chu, G., Liu, J., 2015. New evidence for the presence of Changbaishan Millennium eruption ash in the Longgang volcanic field, Northeast China. *Gondwana Res.* 28, 52–60.
- Tyler, J., Kashiwayama, Y., Ohkouchi, N., Ogawa, N., Yokoyama, Y., Chikaraishi, Y., Staff, R.A., Ikehara, M., Bronk Ramsey, C., Bryant, C., Brock, F., Gotanda, K., Haraguchi, T., Yonenobu, H., Nakagawa, T., 2010. Tracking aquatic change using chlorine-specific carbon and nitrogen isotopes: the last glacial-interglacial transition at Lake Suigetsu, Japan. 2010. *Geochem. Geophys. Geosystems* 11, 9.
- Wada, K., Nakamura, M., Okuno, M., 2001. Identification of source volcano from the chemical compositions of glasses from the widespread ashes in the surface layers of Asahidake volcano, Central Hokkaido, Japan. *Rep. Taisetsuzan Inst. Sci.* 35, 9–18 (In Japanese).
- Zou, H., Fan, Q., Zhang, H., 2010. Rapid development of the great millennium eruption of Changbaishan (Tianchi) volcano, China/North Korea: evidence from U-Th zircon dating. *Lithos* 119, 289–296.